

Amoxapine

(pronounced amox - a - peen)

Why has amoxapine been prescribed?

Amoxapine is an antidepressant. These medications are prescribed to treat depression and other mood disorders such as anxiety. Antidepressants are effective in reducing the severity of the symptoms of depression. They also shorten the time to recovery and prevent depression returning.

Are there any precautions with amoxapine?

Amoxapine is suitable for most people. As with all medicines, however, there are precautions. Your prescriber will usually check that it is safe to prescribe amoxapine, but let your prescriber know if any of the following apply to you, as extra care may be needed:

- a) if you have epilepsy, diabetes, glaucoma, phaeochromocytoma, porphyria or suffer from heart, liver, kidney, thyroid or prostate trouble.
- b) if you are taking any other medication especially anticoagulants such as warfarin, medication for Parkinson's disease, medication for schizophrenia, medication for anxiety or insomnia, medication for high blood pressure or irregular heart beat, amiodarone, cimetidine, HIV medications, anticonvulsants, antidepressants or strong pain killers. Also, tell your prescriber if you buy any medicine 'over the counter' from a pharmacy or supermarket. Please also tell your prescriber if you take any alternative or complementary medicine such as Chinese herbal medicines. St John's wort should not be taken by patients prescribed amoxapine.
- c) if you are pregnant, breast feeding, or wish to become pregnant.

What is the usual dose of amoxapine?

The starting dose is usually 100mg to 150mg in divided doses or as a single dose at night. The dose is then usually slowly increased. The usual dose of amoxapine may be between 100mg and 300mg daily. It may take some time to get to the dose that is most suitable.

How should amoxapine be taken?

Amoxapine can be taken in divided doses or as single dose at night. However, you and your prescriber may decide that it is better for you to take the medicine at another time.

What form does amoxapine come in?

Amoxapine is available only as tablets. The tablets are available as 50mg and 100mg.

What should be done if a dose is missed?

If you forget a dose, take it as soon as you remember. If you miss a whole day's dose – just carry on as normal with the usual dose. If you miss more than a day's dose, speak to your prescriber.

What will happen when amoxapine is first taken?

Amoxapine, like many medicines, does not work straightaway. For example, it may take several days or even weeks for some symptoms to improve. To begin with, some people find that amoxapine may help them feel more calm and relaxed. Later, (usually in two to three weeks) other symptoms should begin to improve. Unfortunately, some people get side effects before starting to feel better. However, most side effects will lessen and should go away after a few weeks. The table below lists some of the side effects associated with amoxapine. There are other possible side effects – we have listed only the most important ones. Ask your doctor, pharmacist or nurse if you are worried about anything else that you think might be a side effect. Further information on side effects is available in the official manufacturer's leaflet.

Reporting side effects

The 'Yellow Card' scheme encourages patients to report any side effects that they feel may be caused by their medication. Ask your doctor, pharmacist or nurse for the forms if you wish to report any side effects. Alternatively, telephone 0800 100 3352 or go to www.mhra.gov.uk.

Summary of side effects				
Side effect	How common is it?	What can be done about it?		
Drowsiness	Common	This should wear off within a few days or weeks. Discuss with your prescriber if this persists. Taking the medication at night-time may help, but beware of drowsiness in the following morning.		
Blurred vision	Common	This should wear off within a few days. Discuss with your prescriber if this persists. It may be necessary to switch to another medication. Symptoms will go away when switched to another medication.		
Dry mouth	Common	This should wear off within a few days. Sugar-free boiled sweets, chewing gum or eating citrus fruits may help. Discuss with the prescriber if this persists.		
Constipation	Common	Eat lots of fibre – fruit and vegetables are good sources. Drink plenty of fluids. Your prescriber may prescribe a laxative.		
Postural hypotension (low blood pressure)	Common	Try not to stand up too quickly. If you feel dizzy do not drive. This dizziness is not dangerous and should wear off after a few days.		
Sexual dysfunction - reduced libido (desire) and lack of orgasm in males and females. Impotence in males	Uncommon	Any changes should be reported to your prescriber. It may be necessary to switch to another medication. Symptoms should go away when switched to another medication.		
Agitation and anxiety	Uncommon	Any changes should be reported to your prescriber.		
Increased appetite and weight gain	Uncommon	Try to eat healthily and take exercise.		
Sweating	Uncommon	Discuss this with your prescriber if it becomes troublesome.		
Tachycardia (palpitations)	Uncommon	Contact your prescriber if you experience palpitations.		
Menstrual changes (women) – periods may stop	Rare	Any changes should be reported to your prescriber.		
Breast growth and milk production (men & women)	Rare	Any changes should be reported to your prescriber.		
Urinary retention (difficulty passing water)	Rare	Contact your prescriber if you have any difficulty passing water.		
<i>Very common = almost everyone affected</i>	<i>Common = many people affected</i>	<i>Uncommon = some people affected</i>	<i>Rare = few people affected</i>	<i>Very rare = very few people affected</i>

Is amoxapine addictive?

Amoxapine is not addictive, but if you stop taking the medicine suddenly, you may experience unpleasant physical feelings. Symptoms are fairly common and include nausea, insomnia, diarrhoea, excess salivation and anxiety. Reducing the dose of amoxapine slowly may make these symptoms less severe.

What about alcohol?

It is recommended that people taking amoxapine should not drink alcohol. This is because both antidepressants and alcohol can cause drowsiness. If the two are taken together, severe drowsiness can result. This can lead to falls and accidents. As well as this, drinking alcohol may make depression and anxiety worse. However, once people are used to taking medication, then very small amounts of alcohol may not be harmful. It pays to be very careful, because alcohol affects people in different ways, especially when they are taking medication. Never drink alcohol and drive. Discuss any concerns you have with your doctor, pharmacist or nurse.

Is it OK to stop taking amoxapine when symptoms go away?

No. If you stop taking amoxapine, your original symptoms are very likely to return. Most people need to remain on amoxapine for at least six months after they have fully recovered. You should always discuss with your prescriber any plans you have to reduce or stop any of your prescribed medicines.

Are there any alternatives to amoxapine?

Yes, there are alternatives available. Although antidepressants have broadly similar therapeutic effects they differ in the side effects they produce. There are more than 20 antidepressants available so most people can find one which suits them. Psychological therapies can also help in some types of depression, either as an alternative to medication or alongside it.



Remember, leaflets like this can only describe some of the effects of medication. You may also find other books or leaflets useful. If you have access to the internet you may find a lot of information there as well, but be careful, as internet based information is not always accurate.

© 2007 United Kingdom Psychiatric Pharmacy Group www.ukppg.org.uk

This leaflet is to help you understand more about your medication. It is not an official manufacturer's Patient Information Leaflet (see emc.medicines.org.uk/) For more information, you could contact your local hospital pharmacy – they might have a helpline – or visit www.nwmhp.nhs.uk/pharmacy.

This leaflet has been supplied by:

Medicines Information
Pharmacy Department
Maudsley Hospital
LONDON SE5 8AZ
TEL: 020 3228 2317

South London and Maudsley **NHS**
NHS Foundation Trust